Hospital	ueensway Carleton Hospital 3045 Baseline Road Nepean, Ontario K2H 8P4		Surname		Given Name		Sex
			Address				
REQUEST FOR: Radiology, Ultrasound,			Dob	D M	Y	Tel. () OHIP - RAMQ	
Ultrasound Radiology Other For an Appointment of	Bone Densitor Mammography Peripheral Dop		W. Name of e		'	OTHI - NAME	
Telephone: 721-4711		-4771	B. Date of ac				
Previous Exam.	Yes No		Social insi	urance No.		Claim No.:	
REGIONS TO BE EXAMINED	FOR ULTRASOUN	D,INDICATE SP	ECIFIC ORGANS		ιт []	RT 🗌	
PERTINENT CLINICAL INFOR For I.V.P. and C.T. if pt. has ren			:				
PATIENT PREGNANT?	DIABETIC	?	ALLERGIES?				
Copy report to Dr.		Signature			(M.D.)	Tel:	

RADI 425-98-09

ADULT PREPARATION

PATIENTS UNDER 16 YEARS CONSULT REFERRING PHYSICIAN RE LAXATIVES

Telepaque is supplied by the X-ray Department

ROY VAC is a special bowel cleansing kit available from most drug stores.

DULCOLAX (Bisacody) is a special laxative available from drug stores.

G.I.S. or SMALL BOWEL SEE THROUGH STUDY

Do not eat or drink after midnight. Allow for at least 2 hours in the Department for Small Bowel See Through Studies.

IVP

Nothing to eat or drink after midnight. Schedule at least one week after barium studies.

BARIUM ENEMA

Follow instructions in the ROY VAC . **Except DO NOT eat or drink after midnight.**

AIR CONTRAST BARIUM ENEMA

3 Day Diet Preparation - Instructions available from the X-ray Department.

DAY BEFORE EXAMINATION - Follow instructions in the ROY VAC .

Except DO NOT eat or drink after midnight.

ULTRASOUND - ABDOMEN (GALLBLADDER, KIDNEYS, etc...)

6 hours prior to examination - nothing further to eat or drink until examination completed.

ULTRASOUND - PELVIS (BOTH MALE AND FEMALE PATIENTS)

Arrive with full bladder. Drink at least 3 - 4 (8oz. size) glasses of fluid (preferably water **BUT NOT** milk or other dairy products) 1 hour prior to the scheduled examination time.

ULTRASOUND - COMBINED ABDOMEN & PELVIS

6 hours prior to examination - nothing further to eat.

Arrive with a full bladder - One hour before the scheduled examination time, drink 3 - 4 (8 oz. size) glasses of fluid (preferably water **BUT NOT** milk or other dairy products).

ULTRASOUND - PROSTATE (TRUSP) Light breakfast only. Take fleet enema 2 hrs, before your appointment. Empty your bladder before the test. Obtain a fleet enema from your pharmacy.

MAMMOGRAPHY

Do not apply any underarm deodorant or talc to upper body before mammography examination:

BONE DENSITOMETRY

Wear loose clothing. No metallic objects (i.e. zipper/snaps/buckles) over left hip and lower back.

For information call Diagnostic Imaging at 721-4700 ext. 2701.